

Chinese Tallowtree



Figure 1. Young Chinese tallowtree and leaf (inset).



Figure 2. Chinese tallowtree with yellow inflorescences.



Figure 3. White fruit on Chinese tallowtree.

Chinese tallowtree [*Triadica sebifera* (L.) Small] [Syn. *Sapium sebiferum* (L.) Roxb.; *Stillingia sebifera* Michx.] is a deciduous to evergreen tree native to China, Japan, and Korea. It was introduced to the United States in the mid- to late 1700s as a seed oil crop. Elliott (1824) reports it naturalized along the coast of South Carolina and Georgia fewer than 100 years after its introduction. It escaped from cultivation in at least nine southern states. This invasive plant tolerates saline to freshwater flooding, shade to full sun, acidic to alkaline soils, and wet to dry soils. It establishes dense, solid stands that crowd out native plant species. Characterized by rapid growth and prolific seed production, it is an extremely competitive invasive plant.

Tallowtree is commonly grown in the United States as an ornamental because of its yellow flowers, red fall foliage, and unusual white fruit, which gave it the name “popcorn tree.” However, Chinese tallowtree is a noxious weed in several states, including Mississippi; it is not, however, considered a federal noxious weed. Once introduced into a landscape and mature enough to produce seeds, it quickly spreads into surrounding landscapes. If allowed to perpetuate, control is difficult and expensive.

Description

Vegetative Growth

Chinese tallowtree generally grows between 30 and 40 feet tall, but it can reach over 50 feet. In the southeastern United States, it is deciduous. Leaves are simple, alternate, and smooth (**Figure 1**). The foliage is bright green in summer but often turns fiery red in the fall.

Flowering

Chinese tallowtree is monoecious, with drooping, yellow tassels of insect-pollinated flowers in the spring (**Figure 2**) followed by white fruit in the fall (**Figure 3**). The white fruit can persist throughout the winter and is thought to be poisonous. Seed production can be heavy, averaging around 100,000 seeds per tree.

Dispersal

Despite being poisonous, fruit is spread by birds and other wildlife. Water also disseminates fruit, as evidenced by seedlings germinating on floodplains. Chinese tallowtree is still cultivated throughout the South, and human dispersal can facilitate invasions in and around residential areas.

Habitat

Chinese tallowtree is an early successional species and often emerges to dominate forested areas and right-of-way clearings. It reduces the number and variety of native species in a location, altering the ecosystem's structure and function. In southern Texas, herbaceous coastal prairies were transformed into closed-canopy tallowtree forests within 10 years. Although they can be damaged by early hard freezes, Chinese tallowtrees tolerate a wide range of environmental conditions.

Distribution

Chinese tallowtree has spread rapidly in the United States. It is widely planted in the southern United States as an ornamental and as a pollen crop for honey production. The full extent of its cold-hardiness is not known, but it seems to be progressing northward.

Control Methods

Chemical

See **Table 1** for recommended chemical treatments. These products can be applied in a variety of methods that include foliar, frill, basal bark, cut stump, and soil applications. **Frill** (also known as hack-and-squirt) requires a hatchet, machete, or ax to create an opening in the bark every 2 inches around the trunk; the herbicide is applied in these openings. **Basal bark** applications are made by spraying the lower 18–24 inches of the entire plant trunk with a bark-penetrating adjuvant. These treatments are most effective on trunks fewer than 4 inches in diameter and when treated just before bud break. **Cut stump** applications are made after the tree has been felled; the woody tissue just inside the bark of the remaining stump is treated to prevent resprouting. All foliar and basal bark applications also require a nonionic surfactant at 32–64 ounces per 100 gallons of spray solution. Imazapyr and hexazinone are absorbed by tree roots, so applications made within two times the dripline of desirable trees may

cause injury or death. When spraying Chinese tallowtrees near water, use herbicide formulations and adjuvants that are labeled for aquatic applications.

Mechanical

Hand removal may be possible for small infestations, although tallowtree stems break easily. Seedlings can be pulled in wet soils if roots are not very deep.

Cultural

Since Chinese tallowtrees tolerate a wide range of environmental conditions and soil types, cultural control is not practical. However, selecting alternative landscape ornamentals will help prevent further infestations.

Table 1. Chemical control for Chinese tallowtree.

Herbicide	Formulation	Method	Rate
Fosamine	4 lb/gallon	Foliar	192 to 768 oz/A or 30% solution
Glyphosate	3 lb ae/gallon	Foliar	1% solution
		Cut stump	undiluted
Hexazinone	2 lb/gallon	Soil	256 to 512 oz/A
Imazapyr	2 lb ae/gallon	Foliar or Soil	32 to 96 oz/A or 2%
		Frill or Cut stump	64 oz+32 oz water
		Foliar	25 oz/A
Imazapyr+ Metsulfuron	72.7%	Foliar	25 oz/A
Triclopyr	4 lb ae/gallon	Foliar	2% solution
		Basal	20% solution+ bark penetrator
		Frill	20% solution
		Cut stump	undiluted

References

- Chinese Tallow Task Force. 2005. Chinese Tallow Management Plan for Florida, Sept. 2005. Florida Exotic Pest Plant Council, Fort Lauderdale, FL 33307.
- Elliott, S. 1824. *A Sketch of the Botany of South Carolina and Georgia in Two Volumes*. Volume II. J. R. Schenck, Charleston, SC.
- USDA, NRCS. 2007. The PLANTS Database (<http://plants.usda.gov>, 5 September 2007). National Plant Data Center, Baton Rouge, LA 70874-4490 USA.

Publication 3189 (POD-02-18)

By **John D. Byrd Jr.**, PhD, Extension/Research Professor, Plant and Soil Sciences; **Victor Maddox**, PhD, Senior Research Associate, Plant and Soil Sciences; and **Randy Westbrook**, PhD, former Invasive Species Specialist, U.S. Geological Survey.



Copyright 2018 by Mississippi State University. All rights reserved. This publication may be copied and distributed without alteration for nonprofit educational purposes provided that credit is given to the Mississippi State University Extension Service.

Produced by Agricultural Communications.

Mississippi State University is an equal opportunity institution. Discrimination in university employment, programs, or activities based on race, color, ethnicity, sex, pregnancy, religion, national origin, disability, age, sexual orientation, genetic information, status as a U.S. veteran, or any other status protected by applicable law is prohibited. Questions about equal opportunity programs or compliance should be directed to the Office of Compliance and Integrity, 56 Morgan Avenue, P.O. 6044, Mississippi State, MS 39762, (662) 325-5839.

Extension Service of Mississippi State University, cooperating with U.S. Department of Agriculture. Published in furtherance of Acts of Congress, May 8 and June 30, 1914. GARY B. JACKSON, Director