

Estrus (Heat) Detection in Cattle



Estrus in cattle is commonly referred to as heat. It occurs every 18 to 24 days in sexually mature, open (nonpregnant) female cattle when they are receptive to mounting activity by bulls or other cows or heifers.

Heat detection is critical to heat synchronization and breeding programs, particularly artificial insemination and embryo transfer programs. Effective heat detection is often the most limiting factor in an artificial insemination program. Heat detection can also be used to monitor onset of puberty in heifers, regularity of estrous cycles in breeding age females, and breeding effectiveness of natural service sires via returns to heat in the cow herd.

Heat detection efficiency (rate) is the percentage of eligible cows seen or detected in heat. Eligible cows are cows eligible for insemination. Heifers have reached puberty if they have resumed normal estrus function (cycling) after calving (typically 40 days or more postcalving), are free of reproductive disorders or reproductive tract infections, and are open. A heat detection rate of 80 to 85 percent should be attainable.

Heat Signs and Detection Methods

Several methods of heat detection can be implemented. Some involve using heat detection aids. Several different methods can be combined to improve heat detection rates and accuracy. These include visual observation, heat mount detectors, tailhead markers (paint, chalk, crayon, paste), chin-ball markers, detector animals, and electronic heat detection devices.

Visual Observation

Visual observation is a commonly used method of heat detection. It involves a trained observer's recognizing

and recording signs of heat. Observable signs of heat include mounting or attempting to mount other cattle, standing to be mounted by other cattle, smelling other females, trailing other females, bellowing, depressed appetite, nervous and excitable behavior, mud on hindquarters and sides of cattle, roughed up tail hair, vulva swelling and reddening, clear vaginal mucous discharge, and mucous smeared on rump.

The surest sign of heat is when a cow or heifer allows other cattle to mount her while she remains standing. This is called standing heat. Cattle may be willing to mount others but may not stand to be mounted when outside of standing heat. This usually indicates she is either coming into or going out of standing heat.

This method requires observation of cattle at least twice daily, typically early in the morning and late in the evening for best results. More frequent observation of cattle for heat improves detection accuracy and increases the likelihood of recognizing the optimal time for breeding cattle, particularly in cattle in which heat is less intense or shorter in duration. Nearly 20 percent more cattle will be observed in heat when checked four times per day versus checking twice daily. Check cattle as often as practical. Space heat detection observation times evenly over 24 hours. Each observation period must be sufficiently long, usually at least 30 minutes, to be effective.

Standing heat can occur any time in a 24-hour period. However, the most likely time for a cow or heifer to show heat signs is at night. The season of the year can influence this, with more cows showing heat at night in hot weather and more showing heat during the day in cold weather. Housing conditions can also

Timeline for Heat Signs in Cattle

	Coming into Heat (8 hours)	Standing Heat (18 hours)	Going out of Heat (14+ hours)
Heat Signs	<ul style="list-style-type: none"> • Stands and bellows • Smells other cows • Headbutts other cows • Attempts to ride other cows but will not stand to be mounted • Red, moist, slightly swollen vulva • Clear mucous discharge from vulva 	<ul style="list-style-type: none"> • Stands to be mounted • Rides other cows • Bellows frequently • Nervous and excitable 	<ul style="list-style-type: none"> • Attempts to ride other cows but will not stand to be mounted • Smells other cows • Clear mucous discharge from vulva

have an effect on the distribution of heat during a 24-hour period. Hot weather, high production, crowded conditions, and high stress environments may reduce mounting activity.

Observers must distinguish among cattle coming into heat, in standing heat, and going out of heat. Females that are in standing heat, were in standing heat yesterday, or will be in standing heat tomorrow are the most likely herd mates to mount other cows or heifers in heat. Observe cows away from the feed bunk so feeding behavior does not interfere with heat detection. Cattle need nonslip footing and ample room to interact freely. Dirt footing increases mounting and standing activity more than concrete footing.

Heat Detection Aids

Heat detection aids are available. They should be used to supplement but not replace visual observation. These include tail paint, Kamar® Heatmount® Detectors, Estrotect™ Heat Detectors, Bovine Beacon®, tail head markers, chin-ball markers, and the HeatWatch® II System, an electronic detection system that records mounting behavior. Heat detection aids differ in their application method, detection method, cost per animal, and detection accuracy.

When Cows Show Heat

Time	Cows showing heat signs, %
6 a.m. to noon	22%
noon to 6 p.m.	10%
6 p.m. to midnight	25%
midnight to 6 a.m.	43%

Source: Cornell University



Cow riding another cow

Detector (teaser) animals can also assist in heat detection. Teaser animals include several types of gomer bulls, which are surgically altered to prevent successful insemination. Select gomer bulls that will not become excessively large. Acceptable disposition and freedom from disease are also important in gomer bulls.

Management Considerations

Good management is important for a successful heat detection program. Animals must have clearly readable, unique identification. An adequate area and equipment for heat detection must be available. This may include binoculars or the ability to approach cattle very closely. Nighttime observations may require artificial lighting, such as security lights, flashlights, or lanterns. Recordkeeping supplies such as paper, writing utensils, and clocks are important, along with a well-organized recordkeeping system.

Persons detecting heat must be well trained in heat detection and recording. Instruct all persons detecting heat to record cow or heifer ID, time of observation, and all signs of heat observed. Record all heat periods detected, even if the cow or heifer will not be bred on that heat. Then breeding wheels, calendars, or heat expectancy charts can be used to help predict future heats.



Cow sniffing the vulva of another cow



Cow standing to be mounted

Heat Detection Aids

Detection Aid	Application ¹	Detection Method	Management Considerations	Approximate Cost
Kamar® Heatmount® Detector	Apply with separate adhesive between tail head and hip bone over the sacrum of cow.	Detector remains white until triggered by weight of mounting animal, then it turns bright red.	Detectors may become partially activated, making it hard to tell if heat has occurred. Detectors can become dislodged from female.	\$1-\$2/head
EstroTECT™ Heat Detector	Apply with self-adhesive between tail head and hip bone over the sacrum of cow or heifer.	Detector remains silver until friction of mounting animal(s) reveals fluorescent color under scratched-off silver layer.	Low branches, gates, and other cattle; can lead to a false positive. Detectors can become dislodged from female.	\$1-\$2/head
Bovine Beacon®	Glue to tail head of cow or heifer.	Contains fluorescent dye that glows in the dark when cow or heifer is mounted by another animal.	Low branches, gates, and other cattle; can lead to a false positive. Detectors can become dislodged from female.	\$1-\$2/head
Tail Head Markers	Smear liberal amounts (at least 2-3" wide) of crayon, chalk, paste, or paint marker on the tail head of cow or heifer.	When marker is rubbed off tail head (hair will be ruffled and pulled back), the cow or heifer has stood to be mounted.	Low branches, gates, other cattle, humidity, and rain can lead to a false positive. May need to reapply every few days.	Less than \$1/head
Chin-Ball Marker	Fit marker device under the chin of a teaser (gomer) bull or androgenized cow or heifer.	An animal wearing the device mounts and slides off the cow or heifer in heat, leaving an ink mark on the back and hip of female.	Maintenance is necessary for continuous use (ink must be refilled). Harness straps may break or stretch. Some markings may be from chin's resting instead of mounting.	\$70-90/harness \$30-40/gallon ink
HeatWatch® II	Place small, digital radio transmitter in a piece of polyester material (patch) and glue onto the tailhead of cow or heifer.	Every time there is a mount, data are sent from that transmitter to a small radio receiver (base station) in the proximity of the heat detection area. Data generated on every mount include the female mounted, date and time of the mount, and duration (in seconds) of each mount.	Patches can become dislodged from females. Transmitters can fall out of patches if not adequately secured. Batteries must have proper charge. Increased heat detection accuracy over other aids.	\$3,950/HeatWatch® II System (base station, access paint, software \$945/repeater \$49/monitor \$3.75/patch \$3.75/10.5-oz caulking tube of glue \$0.028/volatile corrosion inhibitor (VCI) bag

¹Comb application area first to remove dead or shedding hair.

Types of Teaser Animals for Heat Detection

Teaser Animal	Procedure	Advantages	Disadvantages
Penile-Blocked Bull	Veterinarian installs device.	<ul style="list-style-type: none"> • Allows normal mounting. • Prevents extension of penis and breeding. • Helps prevent spread of venereal diseases. 	<ul style="list-style-type: none"> • Effective for only one year, since bulls tend to lose sex drive. • Not frequently available.
Vasectomized Bull	Surgical procedure severing the vas deferens (tubes that carry sperm from testes to penis), causing sterility.	<ul style="list-style-type: none"> • Normal libido • Breeding stimuli by sterile bulls increase conception rate to AI. 	Risk of venereal disease spread.
Prepuce/Penis-Deviated Bull	Redirection of the prepuce and penis; moving the prepuce to one side prevents the penis from lining up with the vulva, thus preventing breeding.	<ul style="list-style-type: none"> • Better for detecting females in heat than either vasectomized or penile-blocked males. • No venereal disease spreading. • No loss of sex drive, thus longer work life. 	In some cases, bulls may be able to copulate.
Caudal-Epididymectomized bull	Removal of epididymis tail to prevent sperm from reaching the penis.	<ul style="list-style-type: none"> • Relatively simple to perform. • Economical method of preparing a teaser bull. 	Risk of venereal disease spread.
Androgenized female	Testosterone propionate injections before the breeding season or Synovex® H implants.	<ul style="list-style-type: none"> • Longer work life. • Safer than bulls. • Injections are cheaper than surgically altering bull. 	Response to treatment can vary.

Summary

The effectiveness of detecting heat in cattle varies, depending on method used. Consider the cost, labor, and management system associated with each method in deciding on the best approach to heat detection. The purpose of heat detection also determines the level of heat detection accuracy needed. For more information on cattle reproduction or related topics, contact an office of the Mississippi State University Extension Service.

References

- Britt, J. H., R. G. Scott, J. D. Armstrong, and M. D. Whitacre. 1986. Determinants of estrous behavior in lactating Holstein cows. *J. Dairy Sci.* 69:2195-2202.
- Hall, J. G., C. Branton, and E. J. Stone. 1959. Estrus, estrous cycles, ovulation time, time of service, and fertility of dairy cattle in Louisiana. *J. Dairy Sci.* 42:1086-1094.
- Hurnik, J. F., King, G. J. and H. A. Robertson. 1975. Estrous and related behaviour in postpartum Holstein cows. *Appl. Anim. Ethol.* 2:55-68.
- Reference to commercial products or trade names within information provided by the Mississippi State University Extension Service does not constitute an endorsement by the Mississippi State University Extension Service and does not imply discrimination against other similar products.

Publication 2610 (POD-01-16)

By Dr. Jane A. Parish, Extension/Research Professor, Animal and Dairy Sciences; Dr. Jamie E. Larson, Associate Professor, Animal and Dairy Sciences; and Dr. Rhonda C. Vann, Research Professor, Brown Loam Branch Experiment Station.



MISSISSIPPI STATE
UNIVERSITY™

EXTENSION

Copyright 2016 by Mississippi State University. All rights reserved. This publication may be copied and distributed without alteration for nonprofit educational purposes provided that credit is given to the Mississippi State University Extension Service.

Produced by Agricultural Communications.

We are an equal opportunity employer, and all qualified applicants will receive consideration for employment without regard to race, color, religion, sex, national origin, disability status, protected veteran status, or any other characteristic protected by law.

Extension Service of Mississippi State University, cooperating with U.S. Department of Agriculture. Published in furtherance of Acts of Congress, May 8 and June 30, 1914. GARY B. JACKSON, Director