

# Plant Analysis Sampling Instructions

This publication tells you how to collect a plant tissue sample for analysis. Remember: Plant analyses and evaluations are worthless unless you submit the proper plant part. Following are instructions as to the plant part, stage of growth, and number of plants to sample. For more information, contact your [county MSU Extension office](#).

## Do Not Sample

- Dead or diseased plants.
- Insect- or mechanically injured plants.
- Stressed plants (those that have suffered from extreme temperatures or moisture).
- Plants with soil-covered leaves.
- Plants in the advanced fruiting stages.
- Plants that have had no rainfall since a foliar application of insecticides, fungicides, or nutrient elements.
- Plants that have shown deficiency symptoms for a prolonged period of time.
- Early in the morning or on extremely cloudy days. Nitrates accumulate under these conditions.

## Do

1. Place plant tissue samples inside a large paper bag. Do not wrap or enclose tissue samples in polyethylene bags (freezer bags) or other impermeable containers. Let succulent or wet tissue samples air-dry for at least one day before mailing to the lab.
2. You may want to compare a normal plant with a suspected nutrient-deficient plant. If so, take two samples—one from the normal plant and the other from the abnormal plant. Place each sample in a separate paper bag. Use individual mailing containers, and make reference to each sample.
3. When sampling instructions are not given for the crop you want to sample, a good rule of thumb is to sample the most recently matured leaves. If you are in doubt, contact your county Extension agent.
4. [Complete the online submission form](#). Once completed, you can drop off or ship your samples to Southern Soil and Plant Lab, LLC, 117 Haley Barbour Parkway, Yazoo City, MS 39194.
5. When possible, collect a soil sample from the same location and time as the plant sample was taken. Send soil samples separately from plant samples, but make reference to each sample provided.

## Be sure to take the proper plant sample:




Aboveground portion of seedlings (less than 12 inches)




Top two or three fully developed leaves at the top of some plants (trifoliates)

Mature leaflet pairs from mature shoots



Top 6 inches of forage

Petioles and leaves from leaves adjacent to fruit



Ear leaf



Soil sample from root area



Table 1. Field Crops

Crop	Stage of Growth	Plant Part to Sample	No. of Plants to Sample
Corn <sup>1</sup>	Seedling stage (less than 12")	All the aboveground portion.	15–20
	Before tasseling	The entire leaf fully developed below the whorl.	10–15
	From tasseling and shooting to silking	The entire leaf at the ear node (or immediately above or below it).	10–15
Soybeans or Other Beans <sup>2</sup>	Seedling stage (less than 12")	All the aboveground portion.	20–30
	Before or during initial flowering	Two or three fully developed leaves at the top of the plant.	20–30
Small Grain (Wheat, Oats, Rye) <sup>3</sup>	Seedling stage (less than 12")	All the aboveground portion.	20–30
	Before heading	The four uppermost leaves.	40–50
Hay, Pasture, or Forage Grasses	Before seed head emergence or at the optimum stage for best quality forage	The four uppermost leaf blades.	40–50
Alfalfa	Before or at 1/10 bloom stage	Mature leaf blades taken about 1/3 of the way down the plant.	40–50
Clover and Other Legumes	Before bloom	Mature leaf blades taken about 1/3 of the way down from the top of the plant.	40–50
Sorghum-Milo	Before or at heading	Second leaf from top of plant.	40–50
Peanuts	Before or at bloom stage	Mature leaves from either the main stem or cotyledon lateral branch.	20–30
Cotton	Before or at first bloom or when first squares appear	Youngest fully mature leaves on main stem.	20–30

<sup>1</sup>Sampling after silking is not recommended.

<sup>2</sup>Sampling after pods begin to set is not recommended.

<sup>3</sup>Sampling after heading is not recommended.

Table 2. Vegetable Crops

Crop	Stage of Growth	Plant Part to Sample	No. of Plants to Sample
Potatoes	Before or during early bloom	Third to sixth leaf from growing tip.	20–30
Head Crops (such as Cabbage)	Before heading	First mature leaves from center of whorl.	20–30
Tomatoes (Field)	Before or during early bloom stage	Third or fourth leaf from growing tip.	20–30
Tomatoes (Greenhouse)	Before or during fruit	Young plants: leaves adjacent to second and third clusters. Older plants: leaves from fourth to sixth clusters.	20–30
Peppers	Before bloom	Most recently matured leaf.	20–30
Beans	Seedling stage (less than 12")	All the aboveground portion.	20–30
	Before or during initial flowering	Two or three fully developed leaves at the top of the plant (trifoliate).	20–30
Root Crops (such as Carrots, Onions, Beets)	Before root or bulb enlargement	Center mature leaves.	20–30
Celery	Midgrowth (12–15" tall)	Petiole of youngest mature leaf.	20–30
Leaf Crops (such as Lettuce, Spinach, Turnips, Mustard)	Midgrowth	Youngest mature leaf.	20–30
Peas	Before or during initial flowering	Leaves from the third node down from the top of the plant.	20–30
Sweet Corn	Before tasseling	The entire fully mature leaf below the whorl.	20–30
	At tasseling	The entire leaf at the ear node.	20–30
Melons (such as Watermelons, Cucumbers, Muskmelons)	Early stages of growth before fruit set	Mature leaves near the base portion of plant on main stem.	20–30

**Table 3. Fruits and Nuts**

Crop	Stage of Growth	Plant Part to Sample	No. of Plants to Sample
Apples, Apricots, Plums, Prunes, Peaches, Pears, Cherries	Midseason	Leaves near base of current year's growth or from spurs.	20–30
Strawberries	Midseason	Youngest full expanded mature leaves.	20–30
Pecans	6–8 weeks after bloom	Leaves from terminal shoots, taking the pairs from the middle of the leaf.	20–30
Walnuts	6–8 weeks after bloom	Mature leaflet pairs from mature shoots.	20–30
Grapes	From end of bloom period through August	Petioles and leaves from leaves adjacent to fruit clusters.	20–30
Berries	Midseason	Youngest mature leaves on laterals or "primo" canes.	20–30

**Table 4. Ornamentals and Flowers**

Crop	Stage of Growth	Plant Part to Sample	No. of Plants to Sample
Ornamental Trees	Current year's growth	Fully developed leaves.	20–30
Ornamental Shrubs	Current year's growth	Fully developed leaves.	20–30
Turf	During normal growing season	Leaf blades. Clip by hand to avoid contamination with soil or other material.	½ pint of material
Roses	During flower production	5-leaflet leaves below bud.	20–30
Chrysanthemums	Before or at flowering	Most recently matured leaf from top of plant.	20–30
Carnations	Unpinched plants	Fourth or fifth leaf pairs from base of plant.	20–30
	Pinched plants	Fifth and sixth leaf pairs from top of primary laterals.	20–30
Poinsettias	Before flowering	Most recently matured, fully expanded leaves.	20–30
Begonias ( <i>Rieger elatior</i> )	Before heavy flower formation	First leaf from top that is 2 or more inches wide.	20–30
Azaleas, Camellias	Before flowering	Most recently matured leaves.	20–30

---

**Publication 1224** (POD-06-26)

Reviewed by **Brendan Zurweller**, PhD, Associate Professor, Plant and Soil Sciences.



Copyright 2026 by Mississippi State University. All rights reserved. This publication may be copied and distributed without alteration for nonprofit educational purposes provided that credit is given to the Mississippi State University Extension Service.

Produced by Agricultural Communications.

Mississippi State University is an equal opportunity institution. Discrimination is prohibited in university employment, programs, or activities based on race, color, ethnicity, sex, pregnancy, religion, national origin, disability, age, sexual orientation, genetic information, status as a U.S. veteran, or any other status to the extent protected by applicable law. Questions about equal opportunity programs or compliance should be directed to the [Office of Civil Rights Compliance](#), 231 Famous Maroon Band Street, P.O. 6044, Mississippi State, MS 39762.

Extension Service of Mississippi State University, cooperating with U.S. Department of Agriculture. Published in furtherance of Acts of Congress, May 8 and June 30, 1914. ANGUS L. CATCHOT JR., Director