

2024 RR2X & XF Soybean Late Maturity Group IV

Variety Response to Iron Deficiency Chlorosis

Brand	Variety	IDC Tolerance Score ¹					Avg. IDC Tolerance Score ²
Gateway Seed	487XFS	5	6	7	7	7	6
Great Heart Seed	GT-4867XF	5	6	7	7	7	6
Innvictis	A4862XF	4	6	7	7	7	6
Innvictis	A4924XF	4	6	7	8	7	6
NK Seeds	NK48-A8XFS	5	6	7	7	7	6
NK Seeds	NK49-N7XF	4	6	7	7	7	6
Progeny	4947XFS	5	6	7	7	7	6
Revere	4826XFS	5	6	7	7	7	6
Armor	48-F55S	6	7	8	8	7	7
Delta Grow	47XF90STS	6	7	8	8	7	7
Delta Grow	48XF33STS	6	7	8	8	8	7
Delta Grow	48XF70STS	5	6	7	7	7	7
Dyna-Gro	S47XF23S	6	7	8	8	7	7
Dyna-Gro	S48XF35	6	7	8	8	8	7
Gateway Seed	473XFS	5	7	8	8	8	7
Gateway Seed	479XFS	6	7	7	7	7	7
Great Heart Seed	GT-4756XF	6	7	7	8	8	7
Great Heart Seed	GT-4791XF	6	7	8	7	7	7
Integra	XF 4875S	6	7	7	7	7	7
Integra	XF4914S	5	7	8	8	8	7
NK Seeds	NK49-C2XFS	5	7	7	8	8	7
Pioneer	P48Z70BLX	5	7	8	8	8	7
Progeny	P 4842XFS	5	7	7	8	7	7
Progeny	P 4848XF	6	7	7	8	8	7
Revere	47-F77	6	7	8	8	7	7
Revere	49-F36	5	7	7	7	7	7
USG	7474XFS	5	7	7	8	8	7
USG	7495XFS	5	7	7	7	7	7

Brand	Variety	IDC Tolerance Score ¹					Avg. IDC Tolerance Score ²
Armor	48-F25S	6	8	8	8	9	8
Delta Grow	48XF80RKN	6	8	9	9	10	8
Delta Grow	49XF85STS	6	8	9	9	9	8
Don Mario Seed	DM48F53	7	8	8	9	8	8
Dyna-Gro	S49XF43S	6	8	8	9	8	8
Great Heart Seed	GT-4762XF	7	9	9	9	9	8
Innvictis	A4814XF	7	8	9	9	9	8
Progeny	4724XFS	7	8	8	8	8	8
Progeny	P 4734XFS	6	8	8	9	9	8
Progeny	P 4798XF	6	8	9	9	9	8
Progeny	P 4806XFS	7	8	8	8	7	8
Progeny	P 4824XF	7	8	9	9	9	8
Revere	48-F72	6	8	9	9	9	8
Asgrow	AG49XF4	7	8	9	9	9	9

¹Tolerance scores were assigned on a scale of 1 to 10, with 1 being completely tolerant and 10 being completely susceptible. The five individual columns under this heading present tolerance scores collected at one-week intervals beginning at 21 days after planting. All scores are displayed as an average from two locations (Monroe and Lowndes Counties, MS).

²Overall tolerance score averaged across all rating intervals and locations ($p < 0.0001$).

These data are intended to serve as an additional resource for variety selection specifically for soils with a history of problems associated with iron deficiency chlorosis. Consult other sources such as results from official variety trials and demonstration programs for detailed information regarding variety performance.

The information given here is for educational purposes only. References to commercial products, trade names, or suppliers are made with the understanding that no endorsement is implied and that no discrimination against other products or suppliers is intended.

Publication 4096 (POD-03-25)

By **Trent Irby**, PhD, Associate Director and Professor, **Brittany Elliott**, Extension Associate II, **William Paul O'Neal**, Extension Technician, **Paul Garrett Oswalt**, Extension Associate II, Plant and Soil Sciences, and **Brad Burgess**, Director, Research Support — Variety Testing.

Copyright 2025 by Mississippi State University. All rights reserved. This publication may be copied and distributed without alteration for nonprofit educational purposes provided that credit is given to the Mississippi State University Extension Service.

Produced by Agricultural Communications.

Mississippi State University is an equal opportunity institution. Discrimination is prohibited in university employment, programs, or activities based on race, color, ethnicity, sex, pregnancy, religion, national origin, disability, age, sexual orientation, genetic information, status as a U.S. veteran, or any other status to the extent protected by applicable law. Questions about equal opportunity programs or compliance should be directed to the [Office of Civil Rights Compliance](#), 231 Famous Maroon Band Street, P.O. 6044, Mississippi State, MS 39762.

Extension Service of Mississippi State University, cooperating with U.S. Department of Agriculture. Published in furtherance of Acts of Congress, May 8 and June 30, 1914. ANGUS L. CATCHOT JR., Director