

Hay Calculator for Livestock Producers Excel Program (V2019) Instruction Sheet

Winter feeding costs make up a large portion of production costs for Mississippi livestock producers. This Excel-based program will help you evaluate hay feeding delivery systems and their economic impacts. It should provide an accurate estimate of the amount of hay needed for winter feeding as well as the necessary supplementation if hay is not meeting the nutrient requirements of a specific livestock class.

It is essential you estimate how much hay you need for a specific feeding period to ensure animal feeding requirements are met. Not providing the necessary levels of crude protein (CP) and total digestible nutrients (TDN) could affect animal performance and may cause economic losses. The hay calculator is based on different classes of livestock produced in Mississippi and will require a hay test to determine the nutritional value of the hay being fed.

The hay calculator is compatible with Excel 93-2007. To download the hay calculator, see Extension <u>Publication P2590</u> <u>Hay Calculator for Livestock Producers, Excel Program (V2019)</u> <u>Instruction Sheet</u> and look for the publication file labeled *hay_ cal.xls*. To use the hay calculator, follow these instructions:

- Make sure the macro function in your computer is enabled. You can change macro security settings in the Trust Center, unless a system administrator in your organization has changed the default settings to prevent you from changing the settings.
- On the Developer tab, in the Code group, click Macro Security. If the Developer tab is not displayed, click the Microsoft Office Button, click Excel Options, and then, in the Popular category, under Top Options for Working with Excel, click Show Developer tab in the Ribbon.
- ▶ In the Macro Settings category, under Macro Settings, click the option you want. Any changes you make in the Macro Settings category in Excel apply only to Excel and do not affect any other Microsoft Office program. You can also access the Trust Center in the Excel Options dialog box. Click the Microsoft Office Button, and then click Excel Options. In the Trust Center category, click Trust Center Settings, and then click the Macro Settings category.

Hay Requirements Calculation

Step 1. Enter the personal information that will help to create a report for record-keeping purposes. Put that information in the yellow highlighted section of the Excel sheet. A reset button at the top left corner of the sheet allows you to reset the values once you have printed the calculations and analysis.

Step 2. A forage quality analysis is required for a more accurate estimate of the amount of nutrients available to meet livestock nutrient requirements. For information related on how to collect a hay sample, please refer to Extension <u>Publication 2539 Hay</u> <u>Testing and Understanding Forage Quality</u>. Send hay samples requiring analysis to the Mississippi State Chemical Laboratory:

Mississippi State Chemical Laboratory P.O. Box CR Mississippi State, MS 39762 Phone: (662) 325-3428 Fax: (662) 325-7807

https://www.mscl.msstate.edu/customers#essential-forms (click Forage Analysis Test Request Form & Limitations Form)

Step 3. You will need to get an inventory of the hay that is available: number of round or square bales and average weight per bale. This helps in determining the amount of hay that is available. It will be beneficial to get the weight of at least 5 to 10 bales within the same lot to estimate the average bale weight in pounds.

Step 4. This step defines how long you plan to feed hay and how much hay loss you will have, based on storage and feeding methods.

- 4.1 Feeding days: How many days you plan to feed hay. This lets you determine if there is a hay surplus or deficiency. Depending on how you store and feed hay, you always have hay losses.
- ► 4.2 Storage losses (Table 1): Estimated values based on type of storage and how long the hay is stored under those conditions gives you values to put into the calculation sheet.
- ► 4.3 Feeding losses: Animals can cause hay losses. Table 2 shows sample ranges in feeding losses.

Table 1. Hay losses under different storage methods.

		% Losses (0–9 months)		% Losses 12–18 months		
Storage	Method	Range	Average	Range	Average	
Outside	Not Covered					
	Ground	5–20	7.5	15–50	17.5	
	Elevated	3–15	6.0	12–35	11.5	
	Covered					
Outside	Ground	5–10	2.5	10–15	7.0	
	Elevated	2–4	2.0	5–10	5.0	
Under roof		2–5	1.5	3–10	3.5	
Barn		0–2	1.0	2–5	2.0	

Source: Huhnkle, R.L., 2003.

Table 2. Hay losses based on feeding system.

	Percent Hay Feeding Losses			
Feeder Type	Range	Average		
Cone	0–4	2		
Cradle	5–15	10		
Ground	10–28	9		
Ring	0–6	3		
Trailer	5–11	6		
Trough	8–20	6		

Source: Buskirk et al., 2003.

Step 5. Select the type(s) of livestock that will be hay-fed on the farm:

- ▶ 5.1. Know the right number of animals that will be fed.
- ▶ 5.2. Know the average weight per animal in pounds to calculate the metabolic weight.
- 5.3. Calculate daily dry matter intake (lb/day) based on the metabolic weight of the animal and adjust based on the crude protein and neutral detergent of the hay.
- 5.4. Hay test report: The percent dry matter "as received." This will let you calculate what percent of that dry matter is moisture and will be used to adjust daily hay fed.
- 5.5. Daily hay fed: Amount of hay needed for each livestock and adjusted for nutrient requirement and losses.
- 5.6. Total hay need: Amount of hay need (in tons) to feed a specific number of animals for a specific time.

▶ 5.7. Nutrient requirement: Determines if the hay meets the nutrient requirements of the livestock. If it doesn't, you will need to supplement. If numbers are in red for crude protein (CP) or total digestible nutrients (TDN), you need to supplement. If numbers are in green, nutrient requirements are met or there is a surplus. **Table 3** provides the nutrient requirements for each livestock class that need to be met with hay or feed supplementation.

Step 6. Hay surplus/deficiency: Based on the total hay needed for a specific feeding time frame. If numbers are red, you need to buy hay. If numbers are green, there is sufficient hay or a surplus.

Step 7. Hay cost analysis lets you calculate how much hay you need, if there is a deficiency of hay, and how much the hay costs. You will need to know the average weight of the bales and the price per bale you will buy in case you need extra hay.

Table 3. Daily nutrient requirements for cattle.

Livestock Type	CP (%)	TDN (%)	Calcium (% ration)	Phosphorus (% ration)	Vitamin A (IU/Ib)
Cattle					
Rep. heifers open	9.6	60.0	0.3	0.2	1300.0
Rep. heifers bred	11.0	65.0	0.4	0.2	1300.0
Lactating 1st calf heifers	12.0	63.0	0.4	0.2	1800.0
Dry 1st calf heifers	9.6	60.0	0.3	0.2	1300.0
Dry brood cow	6.0	45.0	0.2	0.1	1300.0
Stocker steers	13.0	70.0	0.5	0.2	1300.0
Stocker heifers	13.0	70.0	0.5	0.2	1300.0
Breeding bulls	13.0	70.0	0.5	0.2	1800.0
Beef yearling steers, 700–900 lb, medium frame	13.0	70.0	0.5	0.2	1300.0
Beef yearling steers, 800–1000 lb, large frame	12.0	70.0	0.5	0.2	1300.0
Beef yearling heifers, 600-800 lb, medium frame	13.0	70.0	0.5	0.2	1300.0
Beef yearling heifers, 700–900 lb, large frame	12.0	70.0	0.5	0.2	1300.0
Beef cows, 900–1000 lb, moderate milk	11.0	60.0	0.3	0.2	1800.0
Beef cows, 1100–1300 lb, moderate milk	10.7	60.0	0.3	0.2	1800.0
Beef cows, 1300–1500 lb, moderate milk	10.3	59.0	0.3	0.2	1800.0
Beef cows, 900–1000 lb, high milk	13.0	65.0	0.4	0.2	1800.0
Beef cows, 1100–1300 lb, high milk	12.3	63.0	0.4	0.2	1800.0
Beef cows, 1300–1500 lb, high milk	11.8	62.0	0.4	0.2	1800.0
Beef bulls	12.0	70.0	0.5	0.2	1800.0
Dairy cows 1000 lb, 50% forage ration	11.8	62.0	0.4	0.2	1300.0
Horses					
Horses, mature maintenance	8.5	45.0	0.3	0.2	910.0
Horses, mares mid-gestation	9.5	50.0	0.4	0.3	1370.0
Horses, mares late-gestation	11.0	50.0	0.5	0.4	1370.0
Horses, mares 1st 3 months lactation	14.0	70.0	0.5	0.4	1370.0
Horses, mares late-lactation	12.0	60.0	0.4	0.3	1370.0
Horses, weanlings 4–6 months	16.0	50.0	1.0	0.8	910.0
Horses, weanlings 6–12 months	13.5	50.0	0.7	0.6	910.0
Horses, yearlings 12–18 months	12.0	50.0	0.6	0.4	910.0
Horses, 18–24 months	11.0	50.0	0.5	0.4	910.0
Horses, light work	10.0	62.0	0.3	0.2	910.0
Horses, moderate work	10.0	70.0	0.3	0.2	910.0
Horses, heavy work	10.0	80.0	0.3	0.2	910.0
Horses, stallions	11.0	55.0	0.3	0.2	910.0

Livestock Type	CP (%)	TDN (%)	Calcium (% ration)	Phosphorus (% ration)	Vitamin A (IU/lb)	
Sheep						
Mature sheep	9.5	50.0	0.3	0.2	2350.0	
Ewes, winter lamb	11.0	65.0	0.4	0.3	5000.0	
Ewes, May lamb (140% lamp crop)	11.0	65.0	0.4	0.3	5000.0	
Ewes, May lamb (180% lamp crop)	12.0	66.0	0.4	0.3	5000.0	
Ewes, lactating (suckling singles)	12.0	66.0	0.5	0.3	5100.0	
Ewes, lacatating (suckling twins)	13.5	70.0	0.5	0.3	6000.0	
Replacement ewe lambs	12.5	60.0	0.4	0.3	2000.0	
Mature rams	11.0	55.0	0.4	0.3	2500.0	
Goats						
Mature doe with kids	14.0	75.0	0.5	0.4	1400.0	
Weaned kid to yearling	12.0	60.0	0.2	0.2	700.0	
Mature buck	11.0	55.0	0.3	0.2	700.0	
Donkey, 700 lb	6.5	40.0	0.3	0.2	910.0	

Sources: Gimenez, D.M. 1994; NRC, 2000; Kline et al., 2009.

References

- Buskirk, D.D., A.J. Zanella, T.M. Harrigan, J.L. Van Lente, L.M. Gnagey, and M.J. Kaercher. 2003. Large round bale feeder design affects hay utilization and beef cow behavior. J. Anim. Sci. 81: 109-115.
- Gimenez, D.M. 1994. Nutrient requirements of sheep and goats. Alabama Coop. Ext. Serv. ANR-812.
- Huhnke, R.L. 2003. Round bale hay storage. Oklahoma State Coop. Ext. Serv. F-1716.
- Kline, R., S. Porr, and J. Cardina. Horse Nutrition. Ohio State Coop. Ext. Serv. Bulletin 762-00. Online: <u>http://ohioline.osu.</u> <u>edu/b762/index.html</u> (Accessed May 15, 2009).
- Lalman, D. and C. Richards. 2006. Oklahoma State University Feed Commodity Bulletin: Nutrient composition table. Oklahoma State University. Online: <u>http://www.ansi.okstate.</u> <u>edu/exten/feedbull/nutrient.htm</u> (Accessed August 14, 2009).
- National Research Council (NRC). Nutrient requirements for beef cattle. 2000. National Academic Press.

This information is for educational and preliminary planning purposes only. Use this program as a guide only. The hay calculator uses publicly available information as a basis for its calculations; such information may change over time. The user assumes the risk of using or otherwise relying upon any hay calculator output. The hay calculator does not provide supplementation recommendations.

The Mississippi State University Extension Service does not guarantee the functionality of the calculator. MSU Extension does not guarantee the accuracy or completeness of any calculator output. The calculator, its operation, and any output are provided as is and without any express or implied warranty, including merchantability or fitness for a particular purpose. MSU Extension is not bound by any calculator output and is not responsible for use or reliance on such output.

Publication 2590 (POD-11-22)

By Rocky Lemus, PhD, Extension/Research Professor, Plant and Soil Sciences; and Daniel Rivera, PhD, Director of Southwest Research and Extension Center.



Copyright 2022 by Mississippi State University. All rights reserved. This publication may be copied and distributed without alteration for nonprofit educational purposes provided that credit is given to the Mississippi State University Extension Service. Produced by Agricultural Communications.

Mississien i Clata Hairmanita in an anal ann adamita ina

Mississippi State University is an equal opportunity institution. Discrimination in university employment, programs, or activities based on race, color, ethnicity, sex, pregnancy, religion, national origin, disability, age, sexual orientation, gender identity, genetic information, status as a U.S. veteran, or any other status protected by applicable law is prohibited.

Extension Service of Mississippi State University, cooperating with U.S. Department of Agriculture. Published in furtherance of Acts of Congress, May 8 and June 30, 1914. STEVE MARTIN, Interim Director