

# Sex-Sorted Semen for Beef Cow-Calf Production



Attempts to shift the gender ratio for a calf crop have been tried several different ways, but the idea of sorting or selecting semen to favor male or female offspring has been researched since the advent of artificial insemination. This is possible because the fertilizing sperm cell determines the gender of the calf. Because of the genetic makeup of the cells, sperm that produce female offspring are referred to as X-bearing sperm, while those that produce male offspring are referred to as Y-bearing sperm.

## Sex-Sorting Technology

Several attempts have been made to develop a method that efficiently separates bovine semen into fractions containing higher concentrations of X- or Y-bearing sperm. These technologies include sex-specific antibodies, centrifugation, and flow cytometry. Of these attempts, the only method proven to be commercially viable is flow cytometry. This type of sorting was first researched in the 1980s but yielded very low conception rates when sex-sorted semen was used fresh after processing. Work in the laboratory and field studies improved the results, and the first gender-selected calf using frozen semen was produced in 1999. Sexed semen became commercially available on a large scale in the United States in 2004.

The inefficiency of flow cytometry comes from its complexity and slow pace. The principle of this method relies on the fact that X-bearing (female) sperm contain 3.8 percent more DNA than Y-bearing (male) sperm. Before sorting, the sperm cells are stained with a fluorescent dye and then passed through the flow cytometer as drops of liquid containing a single sperm cell per droplet. Because of the difference in amount of DNA, the X-bearing sperm shine brighter than the Y-bearing sperm when exposed to light. This allows the cytometer's laser and detector to determine the gender of the sperm cell based on the amount of light it emits. A positive or negative charge is




Figure 1. Schematic of flow cytometry to sort X-bearing and Y-bearing bovine sperm cells.

then applied to the droplet containing the single sperm cell. Positively charged drops are deflected one way, negatively charged drops are deflected the other, and uncharged droplets pass straight through. The uncharged drops may contain multiple sperm, damaged material, or cells that were not aligned in the proper direction. This process is depicted in Figure 1.

## **Disadvantages of Sex-Sorted Semen**

Sperm cells pass through the flow cytometer at about 60 miles per hour. This seems fast, but, considering they pass in single file and one ejaculate can contain more than seven billion sperm, it takes about three to four times longer to process sex-sorted semen than conventionally processed semen. Therefore, this technique yields fewer straws of frozen semen per ejaculate at an increased cost per straw and results in lower conception rates. Sex-sorted semen is available only on a subset of bulls for which non-sex-sorted semen is available. Under good conditions, pregnancy rates will be about 70 to 90 percent of that for cows or heifers bred with non-sorted semen.

This sorting method is not perfect, but it does shift the ratio to about 85 to 90 percent of the desired sex. As with any other market-driven technology, sex sorting may evolve to become more efficient and less costly. The benefit from shifting the sex ratio of a calf crop has to be weighed against the increased cost and lower fertility.

## **Advantages of Sex-Sorted Semen**

In some regards, sex-sorted semen is more applicable on a large-scale basis to the dairy industry than to the beef industry. Milking herds are almost always more interested in producing replacement heifers, while bull calves are drastically less valuable. With beef cattle herds, the value of shifting the calf crop gender ratio to favor bulls or heifers depends on herd genetics and marketing programs.

For seedstock beef cattle producers, bull sales to commercial cattle producers are often the most significant source of revenue. Heifers are also important for genetic improvement as replacements. For producers who rely on bull sales and require fewer replacement heifers (or purchase bred replacements), breeding with Y-bearing (male) sorted semen could be a practical management decision.

For commercial cattle producers who market feeder calves, steers are usually heavier at weaning and more valuable (\$5.00 to \$10.00 per cwt) than their female counterparts (depending on the method of marketing). Therefore, using Y-bearing sorted semen in a commercial setting would lead to more steer calves that could defray increased semen cost. Furthermore, if heifers are retained

as replacements, individual matings can be hand-selected to produce heifers from more maternal dams and bull calves from terminal crosses. In some beef cattle mating schemes, heifers are more valuable than their male counterparts. A potential example of this is a cross of Brahman and Hereford purebred parents to produce “tiger stripe” calves.

## **Breeding Strategies**

Reserve use of sex-sorted semen for herds where reproductive efficiency has been optimized through intense reproductive management. Pregnancy rates to sexed semen are often highest in virgin heifers that are bred 12 hours after the beginning of standing heat. Mass insemination or timed breeding is not an acceptable breeding method when using sex-sorted semen. Semen handling is similar to traditionally processed semen even though it will likely be packaged in a ¼ cc straw as opposed the usual ½ cc straw. Semen handling during storage and just before insemination is equally critical for success as proper insemination technique. The number of transferable embryos produced by using sex-sorted semen in superovulated cows ranges from one half to 20 percent of that from conventionally packaged semen. Refer to Table 1 for a description of what to expect, relative to conventional semen, when using sex-sorted semen in different production scenarios.

**Table 1. Comparison of Traditional and Sex-Sorted Semen Use for Beef Cattle Operations**

<b>Production consideration</b>	<b>Traditional (non-sex-sorted) semen use</b>	<b>Sex-sorted semen use</b>
Conception rate	Normal conception rate for production situation	Approximately half of the number of transferable embryos using traditional semen
Semen cost	Typical semen cost	Typical semen cost plus premium for sex-sorting technology
Semen availability	Available on a large selection of beef bulls of many breeds	Available on a subset of beef bulls for which traditional semen is available
Resulting calf crop gender ratio	Approximately 50:50 male to female ratio	Approximately 90:10 male to female ratio
Calf crop value	Normal calf crop value for production and marketing situation	Increased calf crop value for gender-selected calves plus decreased calf crop value due to decreased conception rate to sex-sorted semen
Production scenarios where most applicable	No market incentive for significant calf crop gender ratio shift	Market incentives are significant for calf crop gender ratio shift
	Herd reproductive efficiency needs improvement	Herd reproductive efficiency optimized through intense management
	Timed breeding or mass insemination programs	Artificial insemination programs using individual animal heat detection
	Embryo transfer programs where the number of transferable embryos produced is higher priority than embryo gender	Embryo transfer programs where the number of transferrable embryos produced is lower priority than embryo gender

With new technologies that have some potential for controversy, it is important to understand the underlying science before making a judgment on its ethical implications. Sex-sorted semen is not genetically engineered or modified. It is a natural product that is simply separated into fractions that contain a higher percentage of sperm cells that will produce either male or female offspring. Calves from these matings do not have

an increased risk for death or abnormalities, compared to calves from conventional artificial insemination or natural breeding. Calves produced from sex-sorted sperm are expected to grow and develop normally during gestation and after being calved. For more information on beef cattle reproduction or related topics, contact an office of the Mississippi State University Extension Service.

## References

- Johnson, L. A. 1992. Gender preselection in domestic animals using flow cytometrically sorted sperm. *J. Anim. Sci.* 70:8-18.
- Johnson, L. A., G. R. Welch, and W. Rens. 1999. The Beltsville sperm sexing technology: high-speed sperm sorting gives improved sperm output for in vitro fertilization and AI. *J. Anim. Sci.* 77:213-220.
- Tubman, L. M., Z. Brink, T. K. Suh, and G. E. Seidel, Jr. 2004. Characteristics of calves produced with sperm sexed by flow cytometry/cell sorting. *J. Anim. Sci.* 82:1029-1036.

---

Publication 2569 (POD-09-19)

By **Justin D. Rhinehart**, PhD, former Assistant Extension Professor, Animal and Dairy Sciences; and **Jane A. Parish**, PhD, Extension Professor and Head, North MS Research and Extension Center.



*Copyright 2019 by Mississippi State University. All rights reserved. This publication may be copied and distributed without alteration for nonprofit educational purposes provided that credit is given to the Mississippi State University Extension Service.*

Produced by Agricultural Communications.

Mississippi State University is an equal opportunity institution. Discrimination in university employment, programs, or activities based on race, color, ethnicity, sex, pregnancy, religion, national origin, disability, age, sexual orientation, genetic information, status as a U.S. veteran, or any other status protected by applicable law is prohibited. Questions about equal opportunity programs or compliance should be directed to the Office of Compliance and Integrity, 56 Morgan Avenue, P.O. 6044, Mississippi State, MS 39762, (662) 325-5839.

Extension Service of Mississippi State University, cooperating with U.S. Department of Agriculture. Published in furtherance of Acts of Congress, May 8 and June 30, 1914. GARY B. JACKSON, Director