

Beach Vitex (*Vitex rotundifolia* L.f.)



Figure 1. Beach vitex covers a secondary dune on the Isle of Palms near Charleston, SC.



Figure 2. Beach vitex spreads from beachfront properties on Pawleys Island, SC.



Figure 3. Beach vitex leaves and flowers in Point Clear, AL.

Problem

Beach vitex (*Vitex rotundifolia* L.f.) is a deciduous woody vine that was introduced to the Southeastern U.S. from Korea in the mid-1980s. Before its introduction to the southern Atlantic Coast of the U.S., beach vitex had no history of invasiveness. However, by the mid-1990s, dune restoration specialists with the U.S. Army Corps of Engineers began to notice beach vitex spreading from original plantings on South Carolina beaches (Figure 1), crowding out native dune plants, and spreading by seeds and vegetative fragments.

The South Carolina Beach Vitex Task Force has documented beach vitex at more than 115 locations along the South Carolina coast. It also occurs along the North Carolina coast and was recently observed on a beach in Alabama. Efforts are now underway to map and monitor its spread.

Regulations

Currently, beach vitex is not regulated in the U.S. Assessments are now being conducted to determine if the plant should be regulated by federal or state plant regulatory agencies.

Description

Vegetative Growth

Beach vitex can reach 1 to 2 feet high and typically 12 feet long, but runners can grow to infinite lengths, with reports of up to 60 feet (Figure 2). Stems root along their length, forming a mat on the sand. Leaf arrangement is opposite (Figure 3). Leaves are simple, rounded, and gray-green with dense, grayish-white hairs on the lower surface. Leaves are 2 inches long and 1.5 inch wide.

Flowering

Flowers are blue-purple, fragrant, 1 inch wide, and arranged in short inflorescences out of the leaf axils (Figure 3). Fruit are round, ¼-inch wide, and purplish-black when ripe.

Dispersal and Spread

Beach vitex is a prolific seed producer but can also spread from stem (runner) fragments (Figure 2). Seeds float and can be carried in longshore currents and waves to other beaches. Brittle stems can break off during high tides, float away, root, and colonize other beaches. In 2004, free-living plants of beach vitex were detected on North Island, a coastal barrier island near Georgetown, South Carolina, that is accessible only by boat.

Seeds and cuttings that have been chipped up and spread as mulch have also started new populations where dispersed. Beach vitex is fast-growing and drought- and salt-tolerant. Beach vitex is still sold and used as an ornamental, which is another means of dispersal.

Habitat

Currently, beach vitex blankets a number of oceanfront dunes in the Carolinas (Figures 1 and 2). Because of its invasive nature, beach vitex crowds out native dune plants such as sea oats (*Uniola latifolia* L.), American beachgrass (*Ammophila breviligulata* Fern.), and seaside panicum (*Panicum hemitomon* Schult.). In addition to threatening natural sand dune plant communities, beach vitex degrades endangered loggerhead sea turtle (*Caretta caretta* L.) nesting habitat as well as habitat for the federally threatened plant, seabeach amaranth (*Amaranthus pumilus* Raf.).

Distribution

The genus *Vitex* is not native to the lower 48 states, but several species of the genus are widely planted. Beach vitex is reported to be native to Indo-Malaysia and Hawaii, but it is found from China, Taiwan, and Japan south to Malaysia, India, Sri Lanka, Mauritius, Australia, the Pacific Islands, and Hawaii. It has escaped from cultivation in the Southeastern U.S., with free-living populations in Alabama, South Carolina, North Carolina, and Florida (suspected).

In the Midsouth, beach vitex has escaped in Alabama, and mapping efforts are underway to determine if it exists in other Midsouth states.

Control Methods

Chemical

Very little research has been done to evaluate the effectiveness of herbicides on beach vitex. The herbicides listed in **Table 1** are effective for controlling a wide range of woody broadleaf and some grassy plants. These treatments may also damage other broadleaf or grass plants, including some threatened dune species, so it is important to make careful and selective applications. With all foliar treatments, a nonionic surfactant should be added to the spray solution at 1–2 quarts per 100 gallons spray solution.

Herbicide	Formulation	Method	Rate
fosamine	4 lb/gal	foliar	192–768 oz/A or 30% solution
glyphosate	3 lb ae/gal	foliar	2% solution
hexazinone	2 lb/gal	soil	256–512 oz/A
imazapyr	2 lb ae/gal	foliar or soil	32–96 oz/A or 2% solution
imazapyr + metsulfuron	72.7%	foliar	25 oz/A
metsulfuron	60%	foliar	3 oz/A
triclopyr	4 lb ae/gal	foliar	2% solution

Mechanical

Single plants or scattered populations of beach vitex plants should be mechanically removed from an infested site and properly disposed of to avoid unintentional spread of the plant. Digging or pulling plants manually can be used in areas where herbicide use is not desired. Plants should be removed before fruit ripening to avoid seed dispersal. Because stems can propagate through fragmentation, be careful to remove all stems and root fragments.

Physical

No physical control methods are widely used in the United States.

References

- Beach vitex: South Carolina's newest menace. 2007. South Carolina Beach Vitex Task Force. <http://northinlet.sc.edu/resource/vitex.htm#Vitex%20Intro>.
- USDA, NRCS. 2007. The PLANTS Database (<http://plants.usda.gov>, 15 November 2007). National Plant Data Center, Baton Rouge, LA 70874-4490.

More Information

There are approximately 250 species of vitex ranging from temperate to tropical habitats. Beach vitex is in the mint family. A few vitex species are cultivated as woody ornamentals or for beach stabilization. Those commonly cultivated include lilac chastetree (*Vitex agnus-castus* L.), negundo chastetree (*Vitex negundo* L.), beach vitex, and simpleleaf chastetree (*Vitex trifolia* L.). Lilac chastetree and negundo chastetree are grown as shrubs or small trees. Beach and simpleleaf vitex have been considered a single, highly variable species (*Vitex rotundifolia*). However, simpleleaf vitex is upright and shrubby, while beach vitex is a prostrate, sprawling shrub (**Figure 2**).

Publication 3353 (POD-04-19)

Revised by **John D. Byrd Jr.**, PhD, Extension/Research Professor, and **Victor Maddox**, PhD, Senior Research Associate, Plant and Soil Sciences. Originally written by Maddox; Randy Westbrook, PhD, former Invasive Species Specialist, U.S. Geological Survey; Byrd; and Betsy Brabson, South Carolina Beach Vitex Task Force, Georgetown, SC.



Copyright 2019 by Mississippi State University. All rights reserved. This publication may be copied and distributed without alteration for nonprofit educational purposes provided that credit is given to the Mississippi State University Extension Service.

Produced by Agricultural Communications.

Mississippi State University is an equal opportunity institution. Discrimination in university employment, programs, or activities based on race, color, ethnicity, sex, pregnancy, religion, national origin, disability, age, sexual orientation, genetic information, status as a U.S. veteran, or any other status protected by applicable law is prohibited. Questions about equal opportunity programs or compliance should be directed to the Office of Compliance and Integrity, 56 Morgan Avenue, P.O. 6044, Mississippi State, MS 39762, (662) 325-5839.

Extension Service of Mississippi State University, cooperating with U.S. Department of Agriculture. Published in furtherance of Acts of Congress, May 8 and June 30, 1914. GARY B. JACKSON, Director